

Class - X

Mathematics-Basic (241)

Sample Question Paper 2019-20

Max. Marks: 80


Duration: 3 hrs.

General Instructions:

- a) All questions are compulsory
- b) The question paper consists of 40 questions divided into four sections A, B, C & D.
- c) Section A comprises of 20 questions of 1 mark each. Section B comprises of 6 questions of 2 marks each. Section C comprises of 8 questions of 3 marks each. Section D comprises 6 questions of 4 marks each.
- d) There is no overall choice. However internal choices have been provided in two questions of 1 mark each, two questions of 2 marks each, three questions of 3 marks each and three questions of 4 marks each. You have to attempt only one of the alternatives in all such questions.
- e) Use of calculators is not permitted.


SECTION - A

Q 1- 10 are multiple choice questions. Select the most appropriate answer from the given options.

	<p>(a) $2 \text{ Mean} = 3 \text{ Median} - \text{Mode}$ $\text{Median} - \text{Mean}$</p> <p>(c) $\text{Mode} = 2 \text{ Mean} - 3 \text{ Median}$ $\text{Mode} + \text{Mean}$</p>	<p>(b) $2 \text{ Mode} = 3$ $\text{Median} - \text{Mean}$</p> <p>(d) $3 \text{ Median} = 2$ $\text{Mode} + \text{Mean}$</p>	
3.	<p>In the given figure, if TP and TQ are tangents to a circle with centre O, so that $\angle \text{POQ} = 110^\circ$, then $\angle \text{PTQ}$ is</p> <p>(a) 110° (b) 90° (c) 80° (d) 70°</p>		1
4.	<p>325 can be expressed as a product of its primes as</p> <p>(a) $5^2 \times 7$ (b) $5^2 \times 13$ (c) 5×13^2 (d) $2 \times 3^2 \times 5^2$</p>		1
5.	<p>One card is drawn from a well shuffled deck of 52 cards. The probability that it is black queen is</p> <p>(a) $\frac{1}{26}$ (b) $\frac{1}{13}$ (c) $\frac{1}{52}$ (d) $\frac{2}{13}$</p>		1
6.	<p>The sum of the zeroes of the polynomial $2x^2 - 8x + 6$ is</p> <p>(a) - 3 (b) 3 (c) - 4 (d) 4</p>		1
7.	<p>Which of the following is the decimal expansion of an irrational number</p> <p>(a) 4.561 (b) $0.\overline{12}$ (c) 5.010010001... (d) 6.03</p>		1

8. The following figure shows the graph of $y = p(x)$, where $p(x)$ is a polynomial in variable x . The number of zeroes of the polynomial $p(x)$ is **1**

(a) 1 (b) 2 (c) 3 (d) 4



9. The distance of the point $P (3, -4)$ from the origin is **1**

(a) 7 units (b) 5 units (c) 4 units
(d) 3 units

10. The mid point of the line segment joining the points $(-5, 7)$ and $(-1, 3)$ is **1**

(a) $(-3, 7)$ (b) $(-3, 5)$ (c) $(-1, 5)$
(d) $(5, -3)$

(11 - 15) Fill in the blanks:

11. The point which divides the line segment joining the points $A (0, 5)$ and $B (5, 0)$ internally in the ratio 2:3 is _____ **1**


12. The pair of lines represented by the equations $2x+y+3 = 0$ and $4x+ky+6 = 0$ will be parallel if value of k is _____. **1**


OR


If the quadratic equation $x^2 - 2x + k = 0$ has equal roots, then value of k

	is _____.	
13.	The value of $\sin 60^\circ \cos 30^\circ + \sin 30^\circ \cos 60^\circ$ is _____.	1
14.	Value of $\cos 0^\circ \cdot \cos 30^\circ \cdot \cos 45^\circ \cdot \cos 60^\circ \cdot \cos 90^\circ$ is _____.	1
15.	The sides of two similar triangles are in the ratio 2:3, then the areas of these triangles are in the ratio _____	

(16 - 20) Answer the following :

16.	$\triangle PQR$ is right angled isosceles triangle, right angled at R. Find value of $\sin P$.	1
	OR	
	If $15 \cot A = 8$, then find value of $\operatorname{cosec} A$.	
17.	If area of quadrant of a circle is 38.5 cm^2 then find its diameter (use $\pi = \frac{22}{7}$)	1
18.	A dice is thrown once. Find the probability of getting a prime number.	1
19.	In the given fig. If $DE \parallel BC$ Find EC.	1
	 <p>Diagram: A triangle ABC is shown. A horizontal line DE is drawn through the triangle, parallel to the base BC. The line DE intersects the base BC at point E and the side AB at point D. The length of AD is labeled as 1.5 cm, and the length of AE is labeled as 1 cm. The length of AB is labeled as 3 cm.</p>	

	5. $x^2 - 2x + 1$ (i) How many students wrote cubic polynomial (ii) Divide the polynomial $(x^2 + 2x + 1)$ by $(x + 1)$.	10. $x^4 - 1$	
	SECTION C		
27.	Find the zeroes of the quadratic polynomial $x^2 - 3x - 10$ and verify the relationship between the zeroes and coefficient.	3	
28.	Draw a circle of radius 4 cm. From the point 7 cm away from its centre, construct the pair of tangents to the circle. OR Draw a line segment of length 8 cm and divide it in the ratio 2:3	3	
29.	Following figure depicts a park where two opposite sides are parallel and left and right ends are semi-circular in shape. It has a 7m wide track for walking  Two friends Seema and Meena went to the park. Meena said that area of the track is $4066m^2$. Is she right? Explain.	3	
30.	Prove that $\frac{\cot A - \cos A}{\cot A + \cos A} = \frac{\operatorname{cosec} A - 1}{\operatorname{cosec} A + 1}$ OR Prove that: $\frac{\tan A + \sin A}{\tan A - \sin A} = \frac{\sec A + 1}{\sec A - 1}$	3	

31.	Prove that $5 - \sqrt{3}$ is irrational, given that $\sqrt{3}$ is irrational.	3
	OR	
	An army contingent of 616 members is to march behind an army band of 32 members in a parade. The two groups are to march in the same number of columns. What is the maximum number of columns in which they can march ?	
32.	Prove that the lengths of tangents drawn from an external point to a circle are equal.	3
33.	<p>Read the following passage and answer the questions that follows:</p> <p>In a class room, four students Sita, Gita, Rita and Anita are sitting at A(3,4), B(6,7), C(9,4), D(6,1) respectively. Then a new student Anjali joins the class</p> 	3
	<p>(i) Teacher tells Anjali to sit in the middle of the four students. Find the coordinates of the position where she can sit.</p> <p>(ii) Calculate the distance between Sita and Anita.</p> <p>(iii) Which two students are equidistant from Gita.</p>	<p>1</p> <p>1</p> <p>1</p>

34.	Solve $2x + 3y = 11$ and $x - 2y = -12$ algebraically and hence find the value of 'm' for which $y = mx + 3$.	3
	SECTION D	
35.	Find two consecutive positive integers sum of whose squares is 365.	4
36.	If the sum of first 14 terms of an A.P. is 1050 and its first term is 10, find the 20 th term.	4
	OR	
	The first term of an A.P. is 5, the last term is 45 and sum is 400. Find the number of terms and the common difference.	
37.	As observed from the top of a 75m high light house above the sea level, the angles of depression of two ships are 30° and 45° respectively. If one ship is exactly behind the other on the same side of the light house and in the same straight line, find the distance between the two ships. (use $\sqrt{3} = 1.732$)	4
38.	If a line is drawn parallel to one side of a triangle to intersect the other two sides in distinct points, then prove that the other two sides are divided in the same ratio.	4
	OR	
	State and prove the Pythagoras theorem.	
39.	A copper rod of diameter 1 cm and length 8 cm is drawn in to a wire of length 18 m of uniform thickness. Find the thickness of wire.	4
	Or	

	A metallic sphere of radius 4.2 cm is melted and recast into the shape of a cylinder of radius 6 cm. Find the height of the cylinder.													
40.	<p>The following distribution gives the daily income of 50 workers of a factory</p> <table border="1" data-bbox="383 496 1362 686"> <thead> <tr> <th>Daily income</th><th>400-420</th><th>420-440</th><th>440-460</th><th>460-480</th><th>480-500</th></tr> </thead> <tbody> <tr> <th>Number of workers</th><td>12</td><td>14</td><td>8</td><td>6</td><td>10</td></tr> </tbody> </table> <p>Convert this distribution to less than type of cumulative frequency distribution and draw its ogive.</p>	Daily income	400-420	420-440	440-460	460-480	480-500	Number of workers	12	14	8	6	10	4
Daily income	400-420	420-440	440-460	460-480	480-500									
Number of workers	12	14	8	6	10									

Class - X

Mathematics-Basic (241)

Marking Scheme-SQP 2019-20

Max. Marks: 80

Duration: 3 hrs.

1.	(b) 42	(1)
2.	(a) 2 Mean = 3 Median - Mode	(1)
3.	(d) 70°	(1)
4.	(b) $5^2 \times 13$	(1)
5.	(a) $\frac{1}{26}$	(1)
6.	(d) 4	(1)
7.	(c) 5.010010001...	(1)
8.	(c) 3	(1)
9.	(b) 5 units	(1)
10.	(b) (- 3, 5)	(1)
11.	(2, 3)	(1)
12.	2 OR 1	(1)
13.	1	(1)
14.	0	(1)
15.	4:9	(1)
16.	$\sin P = 1/\sqrt{2}$	(1)

OR

$$\operatorname{cosec} A = 17/15$$

17. Area of quadrant = $\frac{1}{4} \times \frac{22}{7} \times r^2 = 38.5$ (use $\pi = \frac{22}{7}$) (1/2)

$$\Rightarrow r = 7\text{cm}$$

$$\therefore \text{diameter} = 14 \text{ cm}$$
(1/2)

18. $\frac{1}{2}$ 1

19. $\frac{AD}{BD} = \frac{AE}{EC}$ (By B.P.T.) (1/2)

$$\frac{1.5}{3} = \frac{1}{EC}$$

$$\therefore EC = 2 \text{ cm}$$
(1/2)

20. $A_5 = a_1 + 4d = 0$ (1/2)

$$1^2 + 4d = 0$$

$$d = -3$$
(1/2)

SECTION - B

21. $P(\text{Two Head}) = \frac{1}{4}$ (1)
(1)

22. Good bulbs = $25 - 5 = 20$ (1)
 $P(\text{good bulb}) = \frac{20}{25} = \frac{4}{5}$ (1)

OR


Of all those outcomes, the ones for which $a + b = 8$ are:

$2+6, 3+5, 4+4, 5+3, 6+2$ or 5 outcomes.

(1)

$$P = 5/36$$
(1)

23.



(1)

$$\angle OLA = 90^\circ$$

$$\angle OMD = 90^\circ$$

$$\angle OLA = \angle OMD$$

Which are alternate angles, hence $AB \parallel CD$

(1)

24.

$$\text{LHS} = \tan 48^\circ \tan 23^\circ \tan 42^\circ \tan 67^\circ$$

(1)

$$= \cot (90^\circ - 48^\circ) \cot (90^\circ - 23^\circ) \tan 42^\circ \tan 67^\circ$$

(1)

$$= \cot 42^\circ \cot 67^\circ \tan 42^\circ \tan 67^\circ$$

$$= 1$$

OR

$$= \cos 48^\circ \cos 42^\circ - \sin 48^\circ \sin 42^\circ$$

(1)

$$= \sin (90^\circ - 48^\circ) \sin (90^\circ - 42^\circ) - \sin 48^\circ \sin 42^\circ$$

(1)

$$= \sin 42^\circ \sin 48^\circ - \sin 48^\circ \sin 42^\circ = 0$$

(1)

25.

$$r = \frac{7}{2}$$

(1)

$$\text{Area of Circle} = \frac{\pi r^2}{4} = \frac{77}{2} \text{ cm}^2$$

(1)

26.

(i) 3 Students

$$\text{(ii)} \quad \frac{x^2 + 2x + 1}{x + 1}$$

(1)

$$= \frac{(x+1)^2}{x+1} = x + 1$$

(1)

SECTION - C

27.	$x^2 - 3x - 10 = 0$ $x^2 - 5x + 2x - 10 = 0$ $x(x-5) + 2(x-5) = 0$ $(x-5)(x+2) = 0$ $x = 5, -2$ $\text{Sum of the roots} = \frac{-b}{a} = \frac{3}{1}$ $\text{which is same as } 5 - 2 = 3$ $\text{product of the roots} = \frac{c}{a} = -10$ $\text{which is same as } 5 \times (-2) = -10$ <p>Hence verified</p>	(3)
28.	<p>Correct construction of given circle (1) Correct construction of two tangents (2)</p> <p>OR</p> <p>Line of given length (1) Correct position of point which divides the line segment in the given ratio (2)</p>	
29.	$\text{Area of track} = 120 \times 70 + [(35)^2 - (120 \times 56 + (28)^2)]$ $= 120 \times 14 + \frac{22}{7} [(35)^2 - (28)^2]$ $= 1680 + \frac{22}{7} \times 7 \times 63$ $= 1680 + 1386$ $= 3066 \text{ m}^2$	(1) $(1\frac{1}{2})$
	Yes, Meena is wrong. $(\frac{1}{2})$	
30.	$\text{L.H.S.} = \frac{\cot A - \cos A}{\cot A + \cos A} = \frac{\frac{\cos A}{\sin A} - \cos A}{\frac{\cos A}{\sin A} + \cos A}$ $= \frac{\cos A (\frac{1}{\sin A} - 1)}{\cos A (\frac{1}{\sin A} + 1)} = \frac{(\frac{1}{\sin A} - 1)}{(\frac{1}{\sin A} + 1)}$ $= \frac{\cosec A - 1}{\cosec A + 1} = \text{R.H.S}$	(1) (1)

OR

$$\begin{aligned}
 \text{L.H.S.} &= \frac{\tan A + \sin A}{\tan A - \sin A} & (1) \\
 &= \frac{\frac{\sin A}{\cos A} + \sin A}{\frac{\sin A}{\cos A} - \cos A} = \frac{\sin A}{\sin A} \frac{[\sec A + 1]}{[\sec A - 1]} & \left(\frac{1}{2}\right) \\
 &= \text{R.H.S} & (1) \\
 & & (1)
 \end{aligned}$$

31. Let us assume that $5 - \sqrt{3}$ is a rational

We can find co prime a & b ($b \neq 0$) such that

$$5 - \sqrt{3} = \frac{a}{b}$$

Therefore $5 - \frac{a}{b} = \sqrt{3}$

So we get $\frac{5b-a}{b} = \sqrt{3}$

Since a & b are integers, we get $\frac{5b-a}{b}$ is rational, and so $\sqrt{3}$ is rational. But $\sqrt{3}$ is an irrational number

Which contradicts our statement

$\therefore 5 - \sqrt{3}$ is irrational

OR

$$616 = 32 \times 19 + 8$$

$$\Rightarrow r = 8 \neq 0$$

$$32 = 8 \times 4 + 0$$


$$\Rightarrow r = 0$$

(2)

The HCF of 32 and 616 is 8.

(1)

32. (1)

	 <p>In $\triangle OPA$ and $\triangle OPB$ $\angle PAO = \angle PBO$ (each 90°) $OP = OP$ (common) $OA = OB$ (radii of same circle) $\triangle OPA \cong \triangle OPB$ (by RHS congruency axiom) Hence $PA = PB$ (CPCT)</p>	(1) (1)
33.	<p>(i) (6,4)</p> <p>(ii) $\sqrt{(6-3)^2 + (1-4)^2} = 3\sqrt{2}$ units</p> <p>(iii) Sita and Rita</p>	(1) (1) (1)
34.	<p>$2x + 3y = 11$ ----(1)</p> <p>$x - 2y = -12$ ----(2)</p> <p>(2) $\Rightarrow x = 2y - 12$ ----(3)</p> <p>Substitute value of x from (3) in (1), we get</p> $2(2y - 12) + 3y = 11$ $\Rightarrow 4y - 24 + 3y = 11$ $\Rightarrow 7y = 35$ $\Rightarrow y = 5$ <p>Substituting value of $y = 5$ in equation (3), we get</p> $x = 2(5) - 12 = 10 - 12 = -2$ <p>Hence $x = -2, y = 5$ is the required solution</p> <p>Now $5 = -2m + 3$ $\Rightarrow 2m = 3 - 5$ $\Rightarrow 2m = -2$ $m = -1$</p>	(1) (1)
		(1)
SECTION - D		
35.	Let two consecutive positive integers be x and $x + 1$	$(\frac{1}{2})$

	$\therefore x^2 + (x + 1)^2 = 365$ $\Rightarrow x^2 + x - 182 = 0$ $(x + 14)(x - 13) = 0$ $\therefore x = 13$ <p>Hence two consecutive positive integers are 13 and 14</p>	(1 $\frac{1}{2}$) (1) (1)
36.	<p>Let common difference be d</p> $\Rightarrow \frac{14}{2}[2(10) + (n - 1)d] = 1050$ $\Rightarrow d = 10$ $a_{20} = a + 19 d$ $= 10 + 19 (10) = 200$	(2) (2)

OR

$$a=5$$

$$a_n = 45$$

$$S_n = 400$$

$$\Rightarrow \frac{n}{2} (5+45) = 400$$

$$50n = 800$$

$$n = 16$$

(2)


$$\text{also } a_n = 45$$

$$5+15d = 45$$

$$15d=40$$

$$d=8/3$$

(2)

37.		(1)
		(1)
38.	<p>For correct fig</p> <p>$\tan 45^\circ = \frac{75}{CD}$</p> <p>$1 = \frac{75}{CD} \Rightarrow CD = 75$</p> <p>$\tan 30^\circ = \frac{75}{BD}$</p> <p>$\frac{1}{\sqrt{3}} = \frac{75}{BD}$</p> <p>$\Rightarrow BD = 75\sqrt{3}$</p> <p>$\Rightarrow \text{Distance between two ships} = BC = 75(\sqrt{3} - 1)\text{m}$</p> <p>$= 54.9 \text{ m}$</p>	(1)
38.	<p>For correct, Given, To prove, construction and Figure</p> <p>For correct proof</p> <p>OR</p> <p>For correct statement, Given, To prove, Construction and Figure</p>	$(4 \times \frac{1}{2} = 2)$ (2) $(5 \times \frac{1}{2} = 2\frac{1}{2})$

	For correct proof	(1½)												
39.	A.T. Q. $\pi r^2 \times 1800 = \pi \times \frac{1}{2} \times \frac{1}{2} \times 8$ $\Rightarrow r^2 = \frac{1}{900}$ $\Rightarrow r = \frac{1}{30}$ $\therefore \text{Thickness of wire} = \frac{1}{15} \text{ cm}$	(2) (1½) (½)												
	OR													
	$\frac{4}{3} \pi r^3 = \pi R^2 h$ $\frac{4}{3} (4.2)^3 = (6)^2 h$ $\Rightarrow h = \frac{2744}{100}$ $\therefore h = 27.44 \text{ cm}$	(2) (1½) (½)												
40.	<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="text-align: center;">Daily Income</th> <th style="text-align: center;">Number of workers</th> <th style="text-align: center;">Cumulative Frequency</th> </tr> </thead> <tbody> <tr> <td style="text-align: center;">400-420</td> <td style="text-align: center;">12</td> <td style="text-align: center;">12</td> </tr> <tr> <td style="text-align: center;">420-440</td> <td style="text-align: center;">14</td> <td style="text-align: center;">26</td> </tr> <tr> <td style="text-align: center;">440-460</td> <td style="text-align: center;">8</td> <td style="text-align: center;">34</td> </tr> </tbody> </table>	Daily Income	Number of workers	Cumulative Frequency	400-420	12	12	420-440	14	26	440-460	8	34	
Daily Income	Number of workers	Cumulative Frequency												
400-420	12	12												
420-440	14	26												
440-460	8	34												

	460-480	6	40		(2)
	480-500	10	50		(2)

Correct Table
Drawing an ogive with co-ordinates
(420,12), (440,26), (460,34), (480,40), (500,50)